



# DISINFECTING GYM WIPES

Suitable for use on: metal, stainless steel, chrome, glass, vinyl, plastic, floors, walls, childrens toys, appliances.

**KILLS 99.99% OF GERMS • MULTI-SURFACE • BLEACH FREE**

Germicidal, Pseudomonacidal, Tuberculocidal, Fungicidal, Virucidal\*

**KEEP OUT OF REACH OF CHILDREN**

### ACTIVE INGREDIENT:

CITRIC ACID	0.60%
OTHER INGREDIENTS:	99.40%
TOTAL	100.00%

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Product is ready to use. Preclean areas with gross filth and heavy soil prior to disinfection. **TO CLEAN/DEODORIZE:** Unfold premoistened cloth and wipe area to be cleaned for 30 seconds or until clean. **TO DISINFECT:** Unfold premoistened cloth and wipe hard non-porous area or item to be disinfected. Wet all surfaces thoroughly. The surface must remain wet for at least 5 minutes. Wipe or let air dry. No rinsing or wiping is required, except on direct food contact surfaces which require a potable water rinse after treatment. Toys: Use only on hard non-porous surfaces and rinse after treatment. **FUNGICIDAL DIRECTIONS:** This product is effective against Trichophyton mentagrophytes (athlete's foot fungus and a cause of ringworm) on inanimate surfaces in locker rooms, dressing rooms, exercise facilities, and shower and bath areas. Follow disinfection directions. **In the presence of 5% blood serum and a contact time of 5 minutes, is effective against:** **GERMICIDAL:** Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella enterica (formerly Salmonella Choleraesuis), Klebsiella pneumoniae, Listeria monocytogenes, Escherichia coli O157:H7, Staphylococcus aureus - MRSA, Staphylococcus epidermidis - MRSE, Enterococcus faecalis Vancomycin Resistant (VRE) **VIRUCIDAL\*:** Respiratory Syncytial Virus, Rotavirus, Human Immunodeficiency Virus Type 1 (Strain HTLV-III<sub>a</sub>), Herpes Simplex Type 1 Virus, Herpes Simplex Type 2 Virus (Strain G), Vaccinia Virus, Influenza A Virus (Strain Hong Kong), Pandemic 2009 H1N1 Influenza A Virus (formerly called swine flu), Adenovirus Type 2, Feline Calicivirus (surrogate for Norovirus) **FUNGICIDAL:** Trichophyton mentagrophytes. **TUBERCULOCIDAL:** Effective against Mycobacterium bovis (BCG) in 5 minutes at 20°C/68°F on previously cleaned hard non-porous inanimate surfaces. **Special Instructions for Cleaning and Decontaminating Surfaces/Objects Soiled with Blood and Body Fluids Which May Contain HIV-1(AIDS virus).** **PERSONAL PROTECTION:** When handling objects soiled with blood or body fluids use disposable latex gloves, gowns, mask and eye coverings. **CLEANING PROCEDURE:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of the disinfectant. **CONTACT TIME:** Do Not Dilute. Leave precleaned surfaces wet for five minutes using mop, cloth or sponge to mitigate HIV-1 virus. Use a 5 minute contact time to kill all bacteria, viruses\* and fungi listed on the label. **DISPOSAL OF INFECTIOUS MATERIAL:** Blood and other fluids should be autoclaved and disposed of according to Federal, State and Local regulations for infectious waste disposal. **DISPENSER DIRECTIONS:** To start feed, remove large cover. Pull through about one inch. Replace cover. Pull out first towelette and snap off at a 90° angle. Remaining towelettes feed automatically ready for their next use. When through using, keep small center cap closed to prevent moisture loss.

### STORAGE AND DISPOSAL:

Do not leave containers open during storage. No special handling is required. **Towelettes:** Do not reuse towelette. Dispose of used towelette in trash. **Dispenser:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available.

### PRECAUTIONARY STATEMENTS

#### HAZARD TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

**FIRST AID IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a Poison Control Center or doctor for further treatment advice.

EPA REG. NO. 34810-36-90011 EPA EST. NO. 10088-WI-2  
1,200 PRE-MOISTENED TOWELETTES 6" X 8" 7 LBS, 8 OZ.

Manufactured for: Raw Athletics, LLC 718 7th St NW Second Floor Washington, DC 20001



8

62344 00013 5